



Contents lists available at SciVerse ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc

Identification of a sphingosine-sensitive Ca^{2+} channel in the plasma membrane of *Leishmania mexicana*

Gustavo Benaim^{a,b,*}, Yael García-Marchán^a, Claudia Reyes^c, Graciela Uzcanga^a, Katherine Figarella^a

^a Instituto de Estudios Avanzados (IDEA), Caracas, Venezuela

^b Instituto de Biología Experimental, Facultad de Ciencias, Universidad Central de Venezuela, Caracas, Venezuela

^c Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas, Venezuela

ARTICLE INFO

Article history:

Received 4 December 2012

Available online 19 December 2012

Keywords:

Leishmania

Calcium channel

Trypanosomatids

Sphingosine

Ca^{2+}

ABSTRACT

The disruption of the intracellular Ca^{2+} homeostasis of *Leishmania mexicana* represents a major target for the action of drugs, such as amiodarone and miltefosine. However, little is known about the mechanism of Ca^{2+} entry to these cells. Here we show the presence of a Ca^{2+} channel in the plasma membrane of these parasites. This channel has many characteristics similar to the human L-type voltage-gated Ca^{2+} channel. Thus, Ca^{2+} entry is blocked by verapamil, nifedipine and diltiazem while Bay K 8644 opened this channel. However, different to its human counterpart, sphingosine was able to open this channel, while other well known sphingolipids had no effect. This fact could have important pharmacological implications.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Leishmania mexicana, the causative agent of cutaneous and mucocutaneous leishmaniasis is a trypanosomatid. The parasite, after being inoculated in the bloodstream by a sandfly in the form of promastigotes, is taken actively by circulating macrophages, where the parasite transforms to the amastigote form. Then, after reaching a critical number, the host-cell is disrupted, discharging the amastigotes, which in turn invade other macrophages repeating the cycle and thus producing the illness [1].

It has been demonstrated that Ca^{2+} is involved in several functions in these parasites [2], including differentiation and host cell invasion [3,4]. The mechanisms involved in the intracellular Ca^{2+} regulation in these parasites are quite well described [1–3]. They possess a distinctive single mitochondria able to accumulate Ca^{2+} by an electrophoretic uniporter, using as driving force for Ca^{2+} uptake, the mitochondrial electrochemical potential [5]. These parasites also possess acidocalcisomes, interesting acidic organelles able to accumulate large amounts of Ca^{2+} [6]. Acidocalcisomes are very relevant concerning the bioenergetic of these parasites since they accumulate Ca^{2+} in combination with pyrophosphate, which is an energy source, alternative to ATP [2,3]. The endoplasmic reticulum is also involved in Ca^{2+} uptake [2,3]. At the plasma membrane, the parasite is able to extrude Ca^{2+} , due to the presence

of a Ca^{2+} -ATPase [7]. However, little is known concerning the mechanisms of Ca^{2+} entry. It has been recently shown that many drugs of current use against this parasite exert their action through the disruption of its intracellular Ca^{2+} homeostasis. Thus, miltefosine, an alkyl-lysophospholipid of general use against leishmaniasis, is known to exert its leishmanicidal action through the opening of a not yet characterized plasma membrane Ca^{2+} channel [8]. Amiodarone and dronedarone, commonly used antiarrhythmic drugs, are known to strongly affect *L. mexicana* [8,9] and other trypanosomatids [10–12], such as *Trypanosoma cruzi*, the causative agent of Chagas disease. These drugs concern the Ca^{2+} homeostasis, affecting the acidocalcisomes, and also collapsing the electrochemical mitochondrial potential, thus inducing the release of Ca^{2+} to the cytoplasm [9,11,12].

Concerning the mechanisms of Ca^{2+} entry to these parasites, the presence of a Ca^{2+} channel in the plasma membrane should be warranted since these parasites are able to significantly change the intracellular Ca^{2+} content depending on different conditions, for example, during cell invasion [4]. However, information related to the presence of such a Ca^{2+} channel is scarce. It has been reported that arachidonic acid induces an intracellular Ca^{2+} increase in *L. mexicana* [13], but part was due to the release of the cation from intracellular organelles, such as mitochondria and acidocalcisomes, since it was also observed in the absence of extracellular Ca^{2+} [13]. Melittin, on the other hand, is able to induce a Ca^{2+} influx, which appears to depend on the activity of a PLD_2 . Interestingly, based on *Leishmania major* genomic evidences, two genes putatively coding for a protein with similar characteristic to those of an L-type voltage-gated Ca^{2+} channel (VGCC) could be present in

* Corresponding author. Address: Instituto de Estudios Avanzados (IDEA), Carretera Nacional Hoyo de la Puerta, Baruta, 1080 Caracas, Venezuela. Fax: +58 212 9035118.

E-mail address: gbenaim@idea.gob.ve (G. Benaim).

these parasites [14]. In the present work we demonstrate the presence of a Ca^{2+} channel in the plasma membrane of *L. mexicana*, with many similarities with the human VGCC. Accordingly, nifedipine and verapamil, classical inhibitors of this channel, known to affect the growth of *Leishmania sp.* [15], block the parasite Ca^{2+} channel. However, differently from its putative human counterpart, the *L. mexicana* Ca^{2+} channel is opened by the sphingolipid sphingosine (Sph), which is known to be present in these parasites [16,17].

2. Materials and methods

2.1. Chemicals

Sphingosine, ceramide, sphingosine-1-P and ceramide-1-P were purchased from Avanti Polar Lipids Inc. EGTA, digitonin, verapamil, nifedipine, diltiazem were from SIGMA. Fura 2-acetoxymethyl ester (FURA 2-AM) was purchased from Molecular Probes.

2.2. Culture of promastigotes of *L. Mexicana*

Promastigotes of *L. mexicana* were cultured in liver infusion – tryptose (LIT) medium supplemented with 10% inactivated fetal bovine serum under continuous agitation at 29 °C as previously reported [8].

2.3. Intracellular Ca^{2+} measurements

Promastigotes were loaded with the fluorescent ratiometric Ca^{2+} indicator Fura 2, to estimate variations on intracellular Ca^{2+} concentration, as described [9]. Briefly, 2×10^8 parasites were collected by centrifugation at 600g for 2 min and washed twice in PBS buffer plus 1% glucose. Then, the parasites were loaded with Fura 2-AM (6 μM), probenecid (12 μM) and pluronic acid (12 μM) in the same buffer at 29 °C, in the dark under continuous agitation for 4 h. The Fura 2-loaded parasites were washed twice by centrifugation and resuspended in Tyrode buffer. Resuspended parasites were placed in a cuvette under continuous stirring at 29 °C in a PerkinElmer 510 Spectrofluorimeter coupled to a fast-filter device that allows the alternating excitation at Ex 340 nm/380 nm [11]. Thus, the conditions of measurement were Ex 340 nm/380 nm and Em 510 nm, and the results were expressed as the ratio values of the λ Em at the two excitation wavelength.

3. Results

Different sphingolipids, such as ceramide, sphingosine (Sph), ceramide-1-P and sphingosine-1-P, are known to increase the intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) in many human cell lines [18,19]. In order to study the possible effect of these sphingolipids on *L. mexicana*, cultured promastigotes were loaded with the Ca^{2+} indicator Fura 2. In Fig. 1A it can be observed that addition of 20 μM Sph, concentration reported to exert its optimal effect on the $[\text{Ca}^{2+}]_i$ in human cell lines [18], is able to induce a large increase in the $[\text{Ca}^{2+}]_i$ of these parasites. Lower concentrations of Sph also increase the $[\text{Ca}^{2+}]_i$ of these parasites, but to a lesser extent in a dose-dependent manner (results not shown). For this reason we used 20 μM Sph along this work. Interestingly, other sphingolipids, such as ceramide, sphingosine-1-P and ceramide-1-P, at even higher concentrations than those known to increase the $[\text{Ca}^{2+}]_i$ in human cell lines [18,19], were not able to affect the intracellular Ca^{2+} content of these parasites (Fig. 1B). Digitonin was added in these experiments to allow the maximal Ca^{2+} entry to the cells. Since digitonin was able to further induce an increase in the Ca^{2+} fluorescence, the Ca^{2+} channel opened by Sph, similar to the human L-type VGCC, should be able to be inactivated.

Nifedipine, verapamil and diltiazem are the canonical inhibitors of the human L-type VGCC. To test these inhibitors on the effect observed by Sph on the $[\text{Ca}^{2+}]_i$, we added sequentially the different inhibitors prior to the addition of the sphingolipid. In Fig. 2A and B it can be observed that nifedipine and verapamil, at concentration used to block the human VGCC, totally blocked the effect of Sph. Diltiazem (10 μM) also abolished the effect of Sph (result not shown).

A very specific agonist of the human L-type VGCC, BayK 8644, has been widely used for the characterization of its function [20]. In Fig. 2C it can be observed that this agonist was able to substitute Sph. Addition of the sphingolipid after BayK 8644 did not induce a further Ca^{2+} release. Accordingly, if Sph is added before the channel agonist, the latter is without effect (Fig. 2D).

We then studied whether the effect of Sph was due to a Ca^{2+} entrance from the extracellular milieu, or instead, it was consequence of its release from intracellular organelles. In Fig. 3A (Top), we showed that when EGTA was added to sequester Ca^{2+} from the extracellular solution, there was a drop in the fluorescence, probably because of the spontaneous release of some Fura 2 from the parasite. Under this condition, addition of Sph, instead of inducing a $[\text{Ca}^{2+}]_i$ increase, produced a dramatic fall. This result is compatible with the exit of the basal intracellular Ca^{2+} to the outer medium,




Fig. 1. Effect of different sphingolipids on the intracellular Ca^{2+} concentration of *L. mexicana*. Promastigotes of *L. mexicana* were loaded with Fura 2, as explained under “Experimentals”. (A) Sphingosine (20 μM) was added (arrow) directly to the stirring cuvette in the presence of 2 mM CaCl_2 . (B) Sphingosine 1-P (Sph 1-P, 20 μM), Ceramide 1-P (Cer 1-P, 20 μM), Ceramide (Cer, 20 μM), Sphingosine (Sph, 20 μM) and Digitonin (DIG, 30 μM) were added (arrows) directly to the stirring cuvette in the presence of 2 mM CaCl_2 . Traces are representative of at least four independent experiments.

due to the presence of EGTA. This result also demonstrates that the channel opened by Sph is able to permit the Ca^{2+} fluxes in both directions, similar to its human analog. In order to study if verapamil could block this Ca^{2+} release, the channel blocker was added before Sph (Fig. 3B). Under this condition the effect of Sph was unaffected, demonstrating that verapamil can only obstruct the channel when acting from outside, probably acting resembling a stopper.

We then test if addition of Sph directly to the parasite population was able to show any discernible effect. We could observe that after 20 min of the addition of 5 μM of the sphingolipid to promastigotes of *L. mexicana*, the parasites started to take a rounded form (compare Fig. 4B to A). At 10 μM of Sph the effect was more evident (Fig. 4C). Finally, at 20 μM , all the parasites were rounded (Fig. 4D). We also observed that, as the concentration of the sphingolipid was increased, the parasites lost its mobility. Even more, we could observe that parasites changes in the shape and mobility were followed by cell death, as assessed with the use of trypan blue exclusion.

4. Discussion

Ca^{2+} is known to exert important functions in the human parasite *L. mexicana* [1]. However, the mechanism of Ca^{2+} entry in these parasites is unknown. In the present work we show compelling evidences for the presence of a plasma membrane Ca^{2+} channel in *L. mexicana*, with characteristics that resemble the human L-type

VGCC. Thus, the channel is blocked by well-known L-type VGCC antagonists, such as verapamil, nifedipine and diltiazem. Even more, this channel can be opened by BayK 8644, a very specific L-type VGCC agonist.

A very interesting finding obtained in the course of this work was that Sph was able to activate the *L. mexicana* plasma membrane Ca^{2+} channel. This sphingolipid is known to inhibit many human enzymes, such as protein kinase C [21] and several Ca^{2+} -calmodulin dependent enzymes [22]. Concerning Ca^{2+} movements, it has been demonstrated that Sph inhibits the human plasma membrane Ca^{2+} -ATPase, PMCA, [23] and is also able to open a TRP channel [24]. Sph is also known to increase the intracellular Ca^{2+} concentration in several cells, such as Jurkat T lymphocytes [18], by a yet unknown mechanism. However, there is no report of an action of Sph in *Leishmania* parasites. Nevertheless, the presence of this and other sphingolipids in these parasites is well documented [16,17]. Experiments conducted with null-mutants of key enzymes from the sphingolipids pathway in *L. major* indicate that sphingolipids are not relevant, at least to support cell growth of promastigotes in culture [16]. Instead, they seem to be necessary for the differentiation to amastigotes and for the production of acidocalcisomes [17,25]. Origin of these neutral sphingolipids is not entirely known. There are even doubts if they are synthesized by the parasites or salvaged from the mammalian host [17–25].

L-type Ca^{2+} channels are found in many excitable cell types, including muscle, neuronal, and endocrine cells, where they initiate Ca^{2+} -dependent responses such as contraction and secretion.




Fig. 2. Effect of Ca^{2+} channel blockers (Nifedipine and Verapamil) and Ca^{2+} channel agonist (BayK 8644) on the intracellular Ca^{2+} concentration of *L. mexicana*. Promastigotes of *L. mexicana* were loaded with Fura 2, as explained under “Experimental”. (A) Nifedipine (Nifed, 2 μM) and then Sphingosine (Sph, 20 μM) were added (arrow) directly to the stirring cuvette in the presence of 2 mM CaCl_2 . (B) Verapamil (Verap, 4 μM) and then Sphingosine (Sph, 20 μM) were added (arrows) directly to the stirring cuvette in the presence of 2 mM CaCl_2 . (C) BayK 8644 (BayK, 2 μM) and then Sphingosine (Sph, 20 μM) were added (arrows) directly to the stirring cuvette, in the presence of 2 mM CaCl_2 . (D) Sphingosine (Sph, 20 μM) and then BayK 8644 (BayK, 2 μM) were added (arrows) directly to the stirring cuvette in the presence of 2 mM CaCl_2 . Traces are representative of at least four independent experiments.




Fig. 4. Effect of Sphingosine on promastigotes of *L. mexicana*. Sphingosine was added to cultured promastigotes at the indicated concentration, and after 20 min the different photographs were taken using an 100X immersion objective and phase contrast microscopy. A, Control. B, Sph, 5 μ M. C, Sph, 10 μ M. D, Sph, 20 μ M. Bars in images represent 10 μ m.

occupied by a threonine residue, a conservative substitution. The amino acid similarities could explain the leishmanial sensitivity to DHP and phenylalkylamines.

The dramatic and relatively rapid effect of Sph on the shape and mobility, leading to the death of the parasites, was unexpected and deserves a special mention. These is not due merely to the Ca^{2+} increase induced by sphingosine since a rapid Ca^{2+} augment do not induce the parasite death [30]. Nevertheless, as a naturally occurring mechanism, it is conceivable that the production of Sph by the cell in a particular moment should be rapid, local and transitory. This is not the case under the experimental conditions set in this work, since Sph was not removed at any moment. It is not even known if the effect of Sph is triggered by direct binding to the channel or through an enzymatic cascade involving kinases or other protein modulators. These experiments, together with electrophysiological studies (i.e. voltage dependency), remain to be done, in order to further characterize this interesting channel.

Acknowledgments

We thank Dr. Lourdes Plaza for critically reviewing the manuscript. This work was supported by grants from Fondo Nacional de Ciencia, Tecnología e Investigación, Venezuela (FONACIT) (No. 2011000884), and from the Consejo de Desarrollo Científico y Humanístico (C.D.C.H.-U.C.V.), Universidad Central de Venezuela to G.B.

References

- [1] G. Benaim, Intracellular calcium signaling and regulation in leishmania, in: F. Tapia, G. Caceres-Dittmar, M.A. Sanchez (Eds.), Molecular and Immune Mechanism in the Pathogenesis of Cutaneous Leishmaniasis, Cap. 5, R.G. Landes Co., Medical Intelligence Unit, Austin, Texas, 1996, pp. 89–106.
- [2] G. Benaim, C.R. Garcia, Targeting calcium homeostasis as the therapy of Chagas' disease and leishmaniasis, *Trop. Biomed.* 28 (2011) 471–481.
- [3] S.N. Moreno, R. Docampo, Calcium regulation in protozoan parasites, *Curr. Opin. Microbiol.* 6 (2003) 359–364.
- [4] H.-G. Lu, L. Zhong, K.P. Chang, et al., Intracellular Ca^{2+} pool content and signaling, and expression of a calcium pump are linked to virulence in *Leishmania mexicana amazonensis*, *J. Biol. Chem.* 272 (1997) 9464–9473.
- [5] G. Benaim, R. Bermúdez, J. Urbina, Ca^{2+} transport in isolated mitochondrial vesicles from *Leishmania braziliensis* promastigotes, *Mol. Biochem. Parasitol.* 39 (1990) 61–68.
- [6] A.E. Vercesi, S.N.J. Moreno, R. Docampo, $\text{Ca}^{2+}/\text{H}^{+}$ exchange in acidic vacuoles exchange in acidic vacuoles of *Trypanosoma brucei*, *Biochem. J.* 304 (1994) 227–233.
- [7] G. Benaim, P. Romero, A calcium pump in plasma membrane vesicles from *Leishmania braziliensis*, *Biochim. Biophys. Acta.* 1027 (1990) 79–84.
- [8] X. Serrano-Martín, G. Payares, M. DeLuca, et al., Amiodarone and miltefosine synergistically induce parasitological cure of mice infected with *Leishmania mexicana*, *Antimicrob. Agents Chemother.* 53 (2009) 5108–5113.
- [9] X. Serrano-Martín, Y. García-Marchan, A. Fernández, et al., Amiodarone destabilizes the intracellular Ca^{2+} homeostasis and the biosynthesis of sterols in *Leishmania mexicana*, *Antimicrob. Agents Chemother.* 53 (2009) 1403–1410.
- [10] G. Benaim, J.M. Sanders, Y. García-Marchan, et al., Amiodarone has intrinsic anti-*Trypanosoma cruzi* activity and acts synergistically with posaconazol, *J. Med. Chem.* 49 (2006) 892–899.
- [11] G. Benaim, V. Hernandez-Rodriguez, S. Mujica, et al., In vitro anti-*Trypanosoma cruzi* activity of dronedarone, a novel amiodarone derivative with an improved safety profile, *Antimicrob. Agents Chemother.* 56 (2012) 3720–3725.
- [12] G. Benaim, A.E. Paniz Mondolfi, The emerging role of amiodarone and dronedarone in treatment of chronic chagasic cardiomyopathy, *Nat. Rev. Cardiol.* 9 (2012) 605–609.
- [13] R. Catisti, S.A. Uyemura, R. Docampo, et al., Calcium mobilization by arachidonic acid in trypanosomatids, *Mol. Biochem. Parasitol.* 105 (2000) 261–271.
- [14] D.L. Prole, C.W. Taylor, Identification of intracellular and plasma membrane calcium channel homologues in pathogenic parasites, *PLoS One* 6 (2011) e26218.
- [15] A.G. Tempote, N.N. Taniwaki, J.Q. Reimão, Antileishmanial activity and ultrastructural alterations of *Leishmania (L.) chagasi* treated with the calcium channel blocker nimodipine, *Parasitol. Res.* 105 (2009) 499–505.
- [16] K. Zhang, J.M. Pompey, F.F. Hsu, et al., Redirection of sphingolipid metabolism toward de novo synthesis of ethanolamine in *Leishmania*, *EMBO J.* 26 (2007) 1094–1104.
- [17] K. Zhang, S.M. Beverley, Phospholipid and sphingolipid metabolism in *Leishmania*, *Mol. Biochem. Parasitol.* 170 (2010) 55–64.
- [18] C. Colina, A. Flores, H. Rojas, et al., Ceramide increase cytoplasmic Ca^{2+} concentration in Jurkat T cells by liberation of calcium from intracellular stores and activation of a store-operated calcium channel, *Arch. Biochem. Biophys.* 436 (2005) 333–345.
- [19] C. Colina, A. Flores, C. Castillo, et al., Ceramide-1-P induces Ca^{2+} mobilization in Jurkat T cells by elevation of $\text{Ins}(1,4,5)\text{-P}_3$ and activation of a store-operated calcium channel, *Biochem. Biophys. Res. Commun.* 360 (2005) 54–60.
- [20] M. Marom, Y. Hagalili, A. Sebag, et al., Conformational changes induced in voltage-gated calcium channel Cav1.2 by BayK 8644 or FPL64176 modify the kinetics of secretion independently of Ca^{2+} influx, *J. Biol. Chem.* 285 (2010) 6996–7005.

- [21] Y.A. Hannun, C.R. Loomis, A.H. Merrill, et al., Sphingosine inhibition of protein kinase C activity and of phorbol dibutyrate binding in vitro and in human platelets, *J. Biol. Chem.* 261 (1986) 12604–12609.
- [22] A.B. Jefferson, H. Schulman, Sphingosine inhibits calmodulin-dependent enzymes, *J. Biol. Chem.* 263 (1988) 15241–15244.
- [23] C. Colina, V. Cervino, G. Benaim, Ceramide and sphingosine have an antagonistic effect on the plasma membrane Ca^{2+} -ATPase from human erythrocytes, *Biochem. J.* 362 (2002) 247–251.
- [24] C. Grimm, R. Kraft, G. Schultz, et al., Activation of the melastatin-related cation channel TRPM3 by D-erythro-sphingosine, *Mol. Pharmacol.* 67 (2005) 798–805.
- [25] K. Zhang, F.F. Hsu, D.A. Scott, et al., *Leishmania* salvage and remodeling of host sphingolipids in amastigote survival and acidocalcisomes biogenesis, *Mol. Microbiol.* 55 (2005) 1566–1578.
- [26] G.H. Hockerman, B.D. Johnson, M.R. Abbott, et al., Molecular determinants of high affinity phenylalkylamine block of L-type calcium channels in transmembrane segment IIIS6 and the pore region of the alpha1 subunit, *J. Biol. Chem.* 272 (1997) 18759–18765.
- [27] G.H. Hockerman, B.Z. Peterson, E. Sharp, et al., Construction of a high-affinity receptor site for dihydropyridine agonists and antagonists by single amino acid substitutions in a non-L-type Ca^{2+} channel, *Proc. Natl. Acad. Sci. USA* 94 (1997) 14906–14911.
- [28] S. Hering, S. Aczél, R.L. Kraus, et al., Molecular mechanism of use-dependent calcium channel block by phenylalkylamines: role of inactivation, *Proc. Natl. Acad. Sci. USA* 94 (1997) 13323–13328.
- [29] S. Yamaguchi, Y. Okamura, T. Nagao, et al., Serine residue in the IIIS5–S6 linker of the L-type Ca^{2+} channel alpha 1C subunit is the critical determinant of the action of dihydropyridine Ca^{2+} channel agonists, *J. Biol. Chem.* 275 (2000) 41504–41511.
- [30] B.E. Cohen, G. Benaim, M.C. Ruiz, et al., Increased calcium permeability is not responsible for the rapid lethal effects of amphotericin B on *Leishmania sp.*, *FEBS Lett.* 259 (1990) 1236–1247.