

Application of Roy adaptation model on adherence to treatment in patients with heart failure

Aplicación del modelo de adaptación Roy sobre la adherencia al tratamiento en pacientes con insuficiencia cardíaca

Shabnam ShariatPanahi¹ <https://orcid.org/0000-0003-2997-1823>, Mansoureh Ashghali Farahan^{2*} <https://orcid.org/0000-0001-9361-9707>, Forough Rafii³ <https://orcid.org/0000-0002-4310-4796>, Maryam Rassouli⁴ <https://orcid.org/0000-0002-5607-8064>, Fereshteh Javaheri Tehran⁵ <https://orcid.org/0000-0002-5894-9492>

¹PhD Student in Nursing, Nursing Care Research Center (NCRC), School of Nursing and Midwifery, Iran University of Medical Sciences, Rashid yasemi St., Valiasr Ave., Tehran, Iran,

²Associate Professor, Nursing Care Research Center (NCRC), School of Nursing and Midwifery, Iran University of Medical Sciences, Rashid yasemi St., Valiasr Ave., Tehran, Iran,

³Professor, Nursing Care Research Center (NCRC), Iran University of Medical Sciences, Rashid yasemi St., Valiasr Ave., Tehran, Iran,

⁴Associate Professor, Cancer Research Center, Shahid Beheshti University of Medical Sciences, Valiasr Ave., Tehran, Iran,

⁵PhD Student in Nursing, Nursing Care Research Center (NCRC), School of Nursing and Midwifery, Iran University of Medical Sciences, Rashid yasemi St., Valiasr Ave., Tehran, Iran,

*Corresponding Author: Mansoureh Ashghali Farahani, Associate Professor, Nursing Care Research Center (NCRC), School of Nursing and Midwifery, Iran University of Medical Sciences, Rashid yasemi St., Valiasr Ave., Tehran, Iran. Corresponding Author EMAIL: m_negar110@yahoo.com

Recibido: 02/03/2020 Aceptado: 01/04/2020 Publicado: 15/05/2020

128

Abstract

Introduction: Poor adherence to treatment in chronic illnesses is a global problem and it's human, clinical and economic consequences is increasing around the world.

Methods: This study aims to construct of the conceptual-theoretical structure which to explains the interaction of environmental stimuli, the coping mechanisms and adaptation modes responses that influence the adherence to treatment in heart failure patient.

Results: We used the three steps of Fawcett's theoretical derivation method, we illustrate the links between the concepts of Roy adaptation model, the concepts and assumptions of the treatment adherence model in patient with heart failure.

Conclusions: It is expected that the results of this study, based on Roy adaptation model, describes adherence behaviors to treatment regimen in patients with heart failure and provide certain strategies and assessment criteria for facilitating adherence to treatment in those patients, and it also could have results for future studies, so that relations between variables are tested and nursing interventions to improve adherence to treatment will be developed and tested.

Keywords: Treatment Adherence and Compliance, Heart failure, adaptation.

Introduction

Introducción: La mala adherencia al tratamiento en enfermedades crónicas es un problema global y sus consecuencias humanas, clínicas y económicas están aumentando en todo el mundo.

Métodos: Este estudio tiene como objetivo construir la estructura conceptual-teórica que explica la interacción de los estímulos ambientales, los mecanismos de afrontamiento y las respuestas de los modos de adaptación que influyen en la adherencia al tratamiento en pacientes con insuficiencia cardíaca.

Resultados: Utilizamos los tres pasos del método de derivación teórica de Fawcett, ilustramos los vínculos entre los conceptos del modelo de adaptación de Roy, los conceptos y supuestos del modelo de adherencia al tratamiento en pacientes con insuficiencia cardíaca.

Conclusiones: Se espera que los resultados de este estudio, basados en el modelo de adaptación de Roy, describan los comportamientos de adherencia al régimen de tratamiento en pacientes con insuficiencia cardíaca y proporcionen ciertas estrategias y criterios de evaluación para facilitar la adherencia al tratamiento en esos pacientes, y también podría tener resultados para estudios futuros, de modo que se desarrollen y prueben las relaciones entre las variables y las intervenciones de enfermería para mejorar la adherencia al tratamiento.

Palabras clave: Cumplimiento y cumplimiento del tratamiento, insuficiencia cardíaca, adaptación.

Contribution of the Paper

“What is already known about the topic?”

The nurses play an important role in managing Heart Failure and improving patients' adherence to the therapeutic regimen.

Roy Adaptation Model is an effective conceptual model for nurses' practice across all setting including heart failure.

Conceptual models are a practical guide to explaining and understanding phenomena of interest.

Middle range theories allow us to study the phenomenon more specifically and with fewer concepts and examine the relationships between concepts through empirically testable hypothesis.

“What this paper adds”?

The results of this research help identify of the conceptual-theoretical structure which to explains the interaction of environmental stimuli, the coping mechanisms and adaptation mode responses that influence the adherence to treatment in HF patient

The Roy Adaptation Model can be used to better management of HF.

Introduction: Heart failure is a global pandemic, about 26 million people around the world are suffering from this disease¹. HF is associated with periods of recurrence and worsening² that would increase costs for health care system considerably³ and reduce patients' life quality^{4,5}. Mortality, morbidity, rehospitalization and disease progression in these patients can be reduced by self-care behaviors or adherence to treatment⁶⁻¹¹.

Adherence to treatment is defined as a range of medical behaviors and lifestyle recommendations, which is in accordance with agreed recommendations by health care providers¹². Adherence to treatment is a multi-dimensional phenomenon that is determined by socioeconomic factors, health care systems related factors, patient related factors, condition rated factors and treatment related factors¹³. Although adherence to treatment is essential, non-adherence to medication and lifestyle recommendation is common in HF¹⁴⁻¹⁶ and its human, clinical and economic consequences, as a chronic illness are increasing worldwide¹⁷.

There are several interventions to improve adherence to treatment in the HF, but even the majority of effective interventions will not result in a significant improvement in adherence and health consequences¹⁸. This is probably related to the complexity of the determinants of adherence¹⁹ and lack of attention to theories that may explain adherence behaviors²⁰. So that, identification of factors influencing adherence to treatment in patients with HF, based on theories, seems essential²¹.

This study aims to construct of the conceptual- theoretical structure which to explains the interaction of environmen-

tal stimuli, the coping mechanisms and adaptation mode responses that influence the adherence to treatment in HF patient. The present study is a part of a greater project named “Designing and testing of treatment adherence model based on Roy adaptation model (RAM) in patients with HF”, which aims to determine the empirical validity of RAM as a conceptual model for explaining and predicting adherence to treatment in this group of patients, through mixed method study in Iran.

Materials and methods

W

e have reviewed literature on implications of RAM on adherence to treatment regimen, in patients with HF in English databases Medline, Web of Science, Scopus and Persian databases SID, Iranmedex, Magiran between the years 2010-2017. The search was conducted on topics related to medical behaviors (medication adherence) and non-pharmalogical recommendations (lifestyle) in HF (Table 1).

Table 1. Search strategy


<p>Medline (“failure”[tiab] OR “heart failure”[tiab] OR “myocardial failure”[tiab] OR “CHF”[tiab] OR “HF”[tiab] OR “congestive heart”[tiab]) AND (“patient concordance”[tiab] OR “patient cooperation”[tiab] OR “patient adherence”[tiab] OR “patient compliance”[tiab] OR “treatment refusal”[tiab] OR “patient dropouts”[tiab] OR “self-care maintenance”[tiab])</p>
<p>Scopus (TITLE-ABS(“failure”) OR TITLE-ABS(“heart failure”) OR TITLE-ABS(“myocardial failure”) OR TITLE-ABS(“CHF”) OR TITLE-ABS(“HF”) OR TITLE-ABS(“congestive heart”)) AND (TITLE-ABS(“patient concordance”) OR TITLE-ABS(“patient cooperation”) OR TITLE-ABS(“patient adherence”) OR TITLE-ABS(“patient compliance”) OR TITLE-ABS(“treatment refusal”) OR TITLE-ABS(“patient dropouts”) OR TITLE-ABS(“self-care maintenance”))</p>
<p>ISI Web of Science (TS=(“failure”) OR TS=(“heart failure”) OR TS=(“myocardial failure”) OR TS=(“CHF”) OR TS=(“HF”) OR TS=(“congestive heart”)) AND (TS=(“patient concordance”) OR TS=(“patient cooperation”) OR TS=(“patient adherence”) OR TS=(“patient compliance”) OR TS=(“treatment refusal”) OR TS=(“patient dropouts”) OR TS=(“self-care maintenance”))</p>

The search results were reviewed by two researchers based on topics and repetitive articles were deleted while reviewing the topics. Then, abstracts of selected articles were retrieved and considering the inclusion and exclusion criteria, were investigated. In this analysis, studies concerned patients with HF who were 18 years old or older, and in terms of severity of illness in Class I-IV

were classified as NYHA or patients with HF_{rEF}, HF_{pEF}, and HF_{mrEF}. Also, studies that have examined one or more components of adherence to treatment and adherence to treatment is the main consequence of the study, results were reported quantitatively and qualitatively were investigated. Editorials, letters to the editor or summaries of conferences were considered as exclusion criteria and subsequently, the original articles were examined based on their relationship with the purpose of this analysis and related studies were selected. Manual search of related resources was also carried out and eventually, 76 studies were investigated (Figure 1).

In the next stage, full text of the qualified studies was reviewed and data analysis was conducted using directed qualitative content analysis method based on RAM. Directed qualitative content analysis contains three stages, preparation, organization and reporting. In preparation stage, text of each article, as a unit of analysis, was read several times by researcher, to become immersed in data. Then, in organization stage, researchers formed the unconstrained matrix of analysis based on the main concepts of RAM, thereby, new categories could be created. In each study, by reading the text line by line, important phrases were highlighted, to be differentiated with other parts. A code is assigned to each important paragraph or phrase. In the next stage, maintaining the coding and performing continuous comparison, obtained codes were sorted based on differences and similarities. After that, based on a conceptual and logical connection, the possibility of placing these categories in the main categories in the matrix or the formation of the new main categories is considered, so that the model of adherence to treatment in patients with HF was designed based on RAM.

Figure 1. Conceptual- theoretical structure



Process of adherence to treatment in patients with HF: The person with HF as an adaptive system is constantly interacting with its changing environment and uses coping processes or defense mechanisms which maintain adaptation in four modes physical status, disease/treatment-related self-concept, social support, role function status.

Stimuli- Three types of stimuli are given to person with HF. When the disease is diagnosed for the person, and he receives a prescription (initiate or continue medication), he must respond with a variety of behaviors such as receiving the correct medication or, in other words, adherence to the medication^{32,33}. For person with HF, HF disease and recommended treatment regimen are considered as the focal stimuli related to adherence to treatment regimen according to RAM. Residual stimuli are internal and/or external environmental factors of human's system, whose effects are not obvious in the given position²⁹ and if identified, they will be placed on the focal and contextual stimuli category³⁴. For this reason, the residual stimuli were not considered in this study.

Contextual stimuli are all of the environmental stimuli that are not the focus of attention or energy, but help to manage the focal stimuli. For persons with HF, factors such as socio-economic factors, personal factors, physical/psychological stress, abilities and coping strategies as the contextual stimuli are related to adherence to treatment.

Most of studies suggest the relationship between socio-economic factors such as income³⁵⁻³⁷, employment status³⁸, education level^{35,39-46}, and ethnicity/race⁴⁵⁻⁵⁰. In only one study was there no relationship between self-care and family income⁴². Most studies examining the association between race/ethnicity and adherence to treatment show that there are differences in adherence to treatment in racial and ethnic groups⁴⁷⁻⁵⁰. There is less adherence to treatment regimen in employed patients due to the problems in coordinating their daily life and job activities in accordance with their physical conditions and their illness³⁸. From 7 studies, 5 studies^{35,39-42} indicate a relationship between higher educational level and patient's better adherence with the treatment regimen.

Personal factors are factors such as age^{36,38,40,42,43,45,46,48,51-55}, gender^{36,41,42,45,46,48,51,53,56-60} and body mass index^{40,56,57}. Age and gender are both positively and negatively associated with adherence to treatment. Considering that HF is more commonly associated with old age, therefore, several factors such as cognitive status, memory impairments, severity of disease, performance status, presence of comorbidity and multiple medications are considered in relation to old age, and potentially prevent effective decision making for patients with HF for self-care and adher-

ence to treatment⁶¹. On the other hand, familial and occupational prohibition and excessive occupation in young people are factors associated with non-adherence with treatment at young age⁵¹.

Results about the relationship between gender and adherence to treatment are different, for instance, better adherence in women is related to their worries about maintaining their health⁵³. On the other hand, women are more vulnerable to self-care behaviors due to having more experience of negative emotions associated with HF and weaker social and family support⁵⁹ and more perceived obstacles (such as lack of scale to control weight at home)⁴¹. Another reason for lower adherence to treatment regimen in women is associated with various etiologies of HF due to gender. HF in men often develops after myocardial infarction and leads to the patient and healthcare system focusing more attention to the disease. However, HF in women is often due to chronic and asymptomatic blood pressure⁵¹.

Abilities and coping strategies such as knowledge of illness^{48,53,54,62,63}, motivation⁶⁴, health literacy^{45,65-67}, self-efficacy^{42,68-71}, daily routine⁷²⁻⁷⁴ and personality (type D)⁷⁵⁻⁷⁷ are other contextual stimuli related to adherence to the treatment regimen. Health literacy and knowledge of illness are important factors of adherence to the treatment regimen, but some studies suggested that knowledge and health literacy cannot guarantee the adherence and the reason for this lack of relevance is the presence of other influential factors on adherence to the treatment regimen such as skill and motivation^{44,63,78}. For example, some of the motivational factors regarding adherence to exercise, relate to social interactions acquired during exercise or motivational factors related to disease conditions and treatment⁶⁴. The results of systematic review confirm that self-efficacy is strongly associated with better self-care⁷¹. The suitability of the treatment regimen and instructions with daily routine and specific needs of each patient related to better adherence to treatment and self-care behaviors^{72,73}.

Physical/psychological stress such as Duration of illness^{31,39,43,46,48,54,55,79}, complexity of medical regimen such as poly-pharmacy and frequency of dosing^{45,53,55,74,80-82}, forgetfulness^{41,74}, comorbidities^{39,41,77,82-87}, depression^{39,46,71,85,88}, anxiety^{35,52,71,72}, sleep disorders^{55,89}, cognitive impairments^{48,56,71,86,90,91}, rehospitalization and outpatient visits^{36,46,53,82} are another contextual stimuli related to adherence to the treatment.

Depression has potential effects on patients' perceptions of their behavior⁹². Although most studies suggest the relationship between depression and adherence to treatment, results of systematic reviews show inconsistent evidence in this regard, and this inconsistency of results may be due to the differences in methodology and also differences in methods of measuring depression and adherence⁸⁵. Anxiety affects people's willingness and ability to engage in self-care, mild to moderate anxiety can activate

individual in the face of threatening events on an adaptive path. In this level, anxiety is a compensatory process, but severe or long lasting anxiety prevents endeavors to adapt²⁹. Although several studies have investigated the relationship between anxiety and adherence to treatment regimen, nature of anxiety and type of self-care consequences must determine whether anxiety should be considered as a facilitating or preventative factor of self-care⁷¹.

Sleep disorder is associated with increase in sleepiness during the day, which leads to reduction in learning, motivation and decision making for self-care performance⁹³. Cognitive impairment is related to reduction in patients' adherence to treatment, through reduced attention and poor performance⁸⁶.

Several studies have investigated the relationship between adherence to treatment and comorbidity, the greater number of comorbidities related to non-adherence is due to the feeling of deterioration, in addition to the fact that the illness also increases the chance of forgetfulness⁹⁴ and on the other hand, apart from multiple comorbidities, it is difficult to differentiate heart failure symptoms from other comorbid illnesses, and this may lead to inability to integrate self-care guidelines for any comorbid illness and eventually less action will be taken when symptoms occur⁸³. Nonetheless, there are different evidences in this regard; researchers concluded that the number of comorbidities does not cause the difference, but rather it is the type of disease which causes the difference⁴⁶.

In the literature related to heart failure, different results were found in the relationship between the severity of illness and adherence to treatment. For example, the severity of illness is related to lower adherence to self-care behaviors and the possible reason for this relationship is the physical disability associated with the severity of illness, so that, the severity of illness may limit the patient's ability to perform daily life activities and subsequently to disregard the treatment regimen⁹⁵. On the other hand, severity of illness has positive relationship with adherence to treatment. In justifying their findings, the researchers stated that the severity of illness could be a factor in provoking patients to follow the regimen and self-care care behaviors to prevent disease progression³⁹. The result systematic review and meta-analysis shows of that history of hospitalization is related to higher levels of adherence, while, there is an inconsistency of evidence between outpatient visits and adherence to treatment⁴⁶. Results of systematic analysis suggest the harmful effects of mild cognitive impairments on self-care, however, it is not possible to combine and compare the findings, due to the limited body of evidence and the lack of compatibility in the tools used⁹¹.

Behavioral Responses—Behavioral responses are internal or external actions and reactions in specific circumstances that show how an individual is adapting to stimuli. Behavioral responses reflect the ability of coping mechanisms to adapt to changing environments and also acts as feedback and additional input to the system. Behavioral responses can be adaptive or ineffective, can be measurable or subjectively reported. Adaptive responses promote the integrity of the human system in terms of adaptation goals such as survival, growth, reproduction, and mastery. For person with HF, adherence to treatment is defined as an adaptive behavior which expresses the perception of people about the cost-benefit of receiving treatment³³. Adherence to treatment reflects the general answer of adaptive system to environmental stimuli and is reflected in four modes of bio-psycho-socio response (physical status, disease/treatment-related self-concept, role function status and social support) with three integrated adaptation levels (100% adherence) compensatory (relative/partial adherence or adherence with help) and compromised (non-adherence).

Defense Mechanism-As mentioned, individuals as an adaptive system are constantly interacting with the environment, which uses intrinsic and acquired mechanisms – subsystem regulator and cognator- to adapt to the environment. The regulator subsystem consists of neuro-chemical and endocrine components that receive stimuli from the internal and external environment and transmitted to the nervous system, and physical responses are seen by the nurse through the activity of the autonomic nervous system, the endocrine and perceptual process. The perceptual process remains in short-term memory; the physical responses can be given as additional stimuli to the system⁹⁶. The cognator subsystem is composed of four psychosocial pathways such as information/perceptual processing, learning, judgment, and emotion. The two subsystems are internal processes Which are not directly observed, but their actions are reflect within the mode of adaptation³⁰. The relationship between two subsystems is through perception and four modes of adaptation, and also the relationship between these subsystems will be different with the adaptive modes, so that the regulator subsystem is often associated with physiological mode and the cognator subsystem with all four modes⁹⁶ Four adaptive modes are also interrelated. Relationships between the four modes of adaptation occur when the stimuli have more than one mode, when the disruptive behavior occurs in more than one mode becomes the stimulus for another modes²⁸.

When the person with HF is given stimulus (for example, medication is prescribed), the regulator subsystem is activated and its response in physical status can be effective

(improvement of symptoms associated with the disease) or ineffective (worse symptoms associated with the disease).The behavioral response observed is stored through the perception in short-term memory for the next reaction, and as stimulus, feedback is given to the system, then the cognator subsystem is activated and the stimuli respond in two ways: threat or challenge⁹⁷.

When a person with HF recognizes a prescribed regimen (focal stimuli) as a threat to self-concept (for example, drug-related side effects, sodium-restricted diet tasteless or exercise-induced tiredness), then anxiety occurs and use emotional responses such as redefining the situation, avoidance, denial or devaluation of situation. These emotion-centered strategies are considered as ineffective responses, and it prevents using problem-centered strategies (for example, seeking treatment and adherence to a regimen) to respond appropriately to the stimulus⁹⁷. So, though self-concept is preserved, non-adherence occurs⁹⁸.

When the treatment regimen is understood as a challenge, problem-centered strategies such as redefining the situation, finding alternative solutions, weighting solutions based on the advantages and disadvantages, choosing the right solution, and ultimately action are used⁹⁷. Therefore, if person understand that the medication helps to improve shortness of breath, or a low salt diet can reduce the amount of edema (adaptive responses in physiological mode), in that case because the benefits of the recommended regimen (i.e. improving dyspnea following diuretic use) more weight than its disadvantages (i.e. the frequency of urine during the night due to diuretic use), In this way, person try to reduce the disadvantages of treatment regimen by using alternative strategies (i.e. the change hours of diuretic use in the early hours of the day to prevent sleep disorders caused by frequent urination Overnight), in which case people will integrate the treatment regimen into their lives and adhere to the treatment regimen.

It should be noted that the relationship between the threat and the challenge in different situations can be changed, although the threat and the challenge can occur simultaneously, but in each situation one will dominate the other, i.e. the existence of a supportive environment and availability more resources make the situations and stimuli perceived as challenges⁹⁹.

Adaptive Modes - The function of subsystems is the achievement of integrated adaptation levels. For persons with HF, adherence to treatment is an integrated adaptation level. Adaptation levels imitate people's behavior and we observe behavior in four adaptive modes: Physical status, disease/treatment-related self-concept, role function status and social support.

For persons with HF, Common symptoms associated with HF and severity of illness is ineffective responses to basic physiological activity. In some studies, severity of illness is related to reduced adherence to treatment⁴⁷⁻¹⁰⁰ and in some studies, it is associated with increased adherence to treatment^{39,48}.

Self-concept is a set of individual's feelings and beliefs which is formed from his inner perception and understanding of other people's reactions. Perceived emotions of illness and treatment can affect individual's self-concept and ultimately his behavior⁹⁸. So, factors such as spirituality¹⁰¹⁻¹⁰³, perception and belief^{7,38,41,46,71,73,104-106} as dimensions of treatment-related self-concept play an important role in self-care performance. Belief is necessary to change the behavior⁴⁵. Individual perception and belief is related to self-care, but the direction and power of relationship is based on the type of belief and self-care behavior that is evaluated⁷¹. Believing in benefits and disadvantages of treatment, necessity of treatment, individual control, chronic disease and lifelong treatment regimen, are factors associated with adherence to treatment^{73,107}. The inner spirituality in praying and saying prayers appears in guiding actions and outer spirituality is described by religious affiliation and helps to understand self-care¹⁰¹⁻¹⁰³.

In RAM, role is considered as a set of expectations of people from each other²⁸. If an individual experiences problem in his roles, this problem affects his ability to manage and improve health, also, change in health can influence an individual's role function status²⁹. Patients with HF usually suffer from social isolation resulted from lifestyle changes, medication regimen, physical limitations associated with shortness of breath and fatigue¹⁰⁸. Also, due to the moderate to severe problems these patients face in doing daily activities, the ability to perform self-care is usually reduced^{109,110}. Interdependence mode relates to the ability to give and receive love, respect and worthiness. In this mode, some relations such as significant others and support system are considered. For persons with HF, social support was used as the representative of interdependent mode, hence, patients' understanding about the provided support is considered as a consistent and effective response in this regard. There is a considerable amount of evidence which suggests the beneficial effects of social support on adherence to treatment regimen and self-care behaviors in HF^{11,42,48,63,106}.

Social support contributes to successful management of chronic disease; nevertheless, the mechanism of its effect has long remained unanswered, maybe because of the mediating effect of self-care confidence. In other words, social support improves self-care confidence/self-efficacy in patients, which will improve patient's ability to perform self-care^{69,111}. In some studies, the results also indicate a lack of relationship between overall social support and its dimensions with adherence to treatment in patients with HF. Maybe the reason for this lack of relationship is measurement of social support and existence of other factors affecting adherence. For example, evidence about the impact of marital status and living arrangement as a structural dimension of social support on adherence to treatment regimen in people with HF has different results⁴⁵. Researchers in justifying their findings, stated that living with a spouse or other family members does not guarantee adherence behavior, but presence of a family member who

has a dietary restriction of sodium can be a strong source of support for the patient's adherence to regimen¹¹².

Assumptions of the model of Adherence to treatment in patients with heart failure: Human health is a function of internal stimuli and the adaptation of levels. Adherence to treatment is conceptualized as an important aspect of the adaptation of levels, reflecting the overall response of the adaptation system to environmental stimuli. Environmental stimuli are processed through regulator and cognator subsystems. These subsystems are internal processes which are not directly observed, but their actions are reflect within the mode of adaptation. Response modes are interrelated, so that the adaptive or ineffective response in each mode may make changes in several adaptive modes that these changes indicate the level of adherence to treatment. In other words, adherence to treatment as a type of adaptation to chronic disease, is the outcome of observed change in the physical status, disease/treatment-related self-concept, role function status and social support which allows patients to feel well-being.

Conclusions

RAM, focusing on the adaptability of people with different environmental stimuli, is one of the most effective models in caring for patients with chronic diseases and an effective guide for nurses in caring for patients with HF. By using this conceptual model, nurses will be able to play an important role to improve the adaptation and integration of patients, through environmental management. Since medication adherence and other lifestyle recommendations in heart failure is a complex and challenging phenomenon for the patient and the treatment team, therefore, the results of this study, based on the RAM, described the behaviors of adherence to treatment regimen in patients with HF and can provide criteria to facilitate adherence to treatment in HF. It can also have implications for future researches, so that, the relationships between variables are tested and nursing interventions are developed and tested to improve adherence to treatment.

Abbreviations:

EF	Ejection Fraction
HF	heart failure
HFmrEF	HF with mid-range ejection fraction
HFpEF	HF with preserved ejection fraction
HFrfEF	HF with reduced ejection fraction
NYHA Classification	New York Heart Association Classification

Acknowledgment: This study is a part of a PhD project named "Designing and testing of treatment adherence model based on Roy adaptation model in patients with heart failure", approved by Iran University of Medical Sciences under the code 95-04-28-29754. Hereby, we wish to express our gratitude to the vice chancellors of Iran universities.

1. Savarese G, Lund LH. Global public health burden of heart failure. *Cardiac failure review*. 2017;3(1):7.
2. Dunlay SM, Redfield MM, Weston SA, Therneau TM, Long KH, Shah ND, et al. Hospitalizations after heart failure diagnosis: a community perspective. *Journal of the American College of Cardiology*. 2009;54(18):1695-702.
3. Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al. Heart disease and stroke statistics—2018 update: a report from the American Heart Association. *Circulation*. 2018;137(12):e67-e492.
4. Gastelurrutia P, Lupón J, Altimir S, de Antonio M, González B, Cabanes R, et al. Effect of fragility on quality of life in patients with heart failure. *The American journal of cardiology*. 2013;112(11):1785-9.
5. Gonzalez DG. *The Effects of Congestive Heart Failure on Quality of Life: As Evaluated by the Minnesota Living with Heart Failure Questionnaire*. 2016.
6. Evangelista LS, Hamilton MA, Fonarow GC, Dracup K. Is exercise adherence associated with clinical outcomes in patients with advanced heart failure? *The Physician and sportsmedicine*. 2010;38(1):28-36.
7. Heydari A, Ziaee ES, Ebrahimzade S. The frequency of rehospitalization and its contributing factors in patient with cardiovascular diseases hospitalized in selected hospitals in Mashhad in 2010. *The Horizon of Medical Sciences*. 2011;17(2):65-71.
8. Lee CS, Moser DK, Lennie TA, Riegel B. Event-free survival in adults with heart failure who engage in self-care management. *Heart & Lung: The journal of acute and critical care*. 2011;40(1):12-20.
9. Mangla A, Kane J, Beaty E, Richardson D, Powell LH, Calvin Jr JE. Comparison of predictors of heart failure-related hospitalization or death in patients with versus without preserved left ventricular ejection fraction. *The American journal of cardiology*. 2013;112(12):1907-12.
10. van der Wal MH, van Veldhuisen DJ, Veeger NJ, Rutten FH, Jaarsma T. Compliance with non-pharmacological recommendations and outcome in heart failure patients. *European heart journal*. 2010;31(12):1486-93.
11. Wu J-R, Frazier SK, Rayens MK, Lennie TA, Chung ML, Moser DK. Medication adherence, social support, and event-free survival in patients with heart failure. *Health Psychology*. 2013;32(6):637.
12. Sabaté E. *Adherence to long-term therapies: evidence for action*: World Health Organization; 2003.
13. van der Wal MH, Jaarsma T. Adherence in heart failure in the elderly: problem and possible solutions. *International journal of cardiology*. 2008;125(2):203-8.
14. Gandhi J, McCue A, Cole R. Nonadherence in the advanced heart failure population. *Current heart failure reports*. 2016;13(2):77-85.
15. Jaarsma T, Strömberg A, Gal TB, Cameron J, Driscoll A, Duengen H-D, et al. Comparison of self-care behaviors of heart failure patients in 15 countries worldwide. *Patient education and counseling*. 2013;92(1):114-20.
16. Krueger K, Botermann L, Schorr SG, Griese-Mammen N, Laufs U, Schulz M. Age-related medication adherence in patients with chronic heart failure: A systematic literature review. *International journal of cardiology*. 2015;184:728-35.
17. Cutler RL, Fernandez-Llimos F, Frommer M, Benrimoj C, Garcia-Cardenas V. Economic impact of medication non-adherence by disease groups: a systematic review. *BMJ open*. 2018;8(1):e016982.
18. Nieuwlaet R, Wilczynski N, Navarro T, Hobson N, Jeffery R, Keepanaseril A, et al. Interventions for enhancing medication adherence. *The Cochrane Library*. 2014.
19. Molloy GJ, O'carroll RE, Witham MD, McMurdo ME. Interventions to enhance adherence to medications in patients with heart failure: a systematic review. *Circulation: Heart Failure*. 2012;5(1):126-33.
20. Holmes EA, Hughes DA, Morrison VL. Predicting adherence to medications using health psychology theories: a systematic review of 20 years of empirical research. *Value in Health*. 2014;17(8):863-76.
21. Wu J-R, Corley DJ, Lennie TA, Moser DK. Effect of a medication-taking behavior feedback Theory-Based intervention on outcomes in patients with heart failure. *Journal of cardiac failure*. 2012;18(1):1-9.
22. Fawcett J, Desanto-Madeya S. *Contemporary nursing knowledge: Analysis and evaluation of nursing models and theories*: FA Davis; 2012.
23. Fawcett J, Garity J. *Evaluating research for evidence-based nursing practice*: FA Davis; 2008.
24. Patton DE, Hughes CM, Cadogan CA, Ryan CA. Theory-based interventions to improve medication adherence in older adults prescribed polypharmacy: a systematic review. *Drugs & aging*. 2017;34(2):97-113.
25. Leventhal H, Cameron L. Behavioral theories and the problem of compliance. *Patient Education and Counseling*. 1987;10(2):117-38.
26. Munro S, Lewin S, Swart T, Volmink J. A review of health behaviour theories: how useful are these for developing interventions to promote long-term medication adherence for TB and HIV/AIDS? *BMC public health*. 2007;7(1):104.
27. Jennings KM. *The Roy Adaptation Model: A Theoretical Framework for Nurses Providing Care to Individuals With Anorexia Nervosa*. *Advances in Nursing Science*. 2017;40(4):370-83.
28. Alligood MR. *Nursing theorists and their work*: Elsevier Health Sciences; 2014.
29. Roy C. *The Roy adaptation model*. New Jersey: Pearson Education; 2009.
30. Whittmore R, Roy SC. Adapting to diabetes mellitus: a theory synthesis. *Nursing Science Quarterly*. 2002;15(4):311-7.
31. Ursavaş FE, Karayurt Ö, İseri Ö. Nursing approach based on Roy Adaptation Model in a patient undergoing breast conserving surgery for breast cancer. *The journal of breast health*. 2014;10(3):134.
32. Naga BS, Al-Atiyyat NM. The Relationship between Pain Experience and Roy Adaption Model: Application of Theoretical Framework. *Middle East Journal of Nursing*. 2014;101(1149):1-6.
33. Kartzos CR. *The Relationship between Intergenerational Solidarity and Medication Adherence in Community Dwelling Elders*. 2013.
34. Nuamah IF, Cooley ME, Fawcett J, McCorkle R. Testing a theory for health-related quality of life in cancer patients: A structural equation approach. *Research in nursing & health*. 1999;22(3):231-42.
35. Piotrowicz E, Orzechowski P, Chrapowicka A, Piotrowicz R. How should we advise heart failure patients on exercise and what should we tell them? *Current heart failure reports*. 2014;11(3):274-80.
36. Chan DC, Shrank WH, Cutler D, Jan S, Fischer MA, Liu J, et al. Patient, physician, and payment predictors of statin adherence. *Medical care*. 2010;196-202.

37. Mead H, Andres E, Ramos C, Siegel B, Regenstein M. Barriers to effective self-management in cardiac patients: the patient's experience. Patient education and counseling. 2010;79(1):69-76.
38. Mudge AM, Adsett J. Factors Predicting Successful Transition to Community-based Maintenance Exercise Programs Following Exercise Rehabilitation. Cardiopulmonary Physical Therapy Journal. 2013;24(4):18.
39. Beker J, Belachew T, Mekonin A, Hailu E. Predictors of adherence to self-care behaviour among patients with chronic heart failure attending Jimma University Specialized Hospital Chronic Follow up Clinic, South West Ethiopia. Journal of Cardiovascular Diseases & Diagnosis. 2014.
40. Juarez DT, Williams AE, Chen C, Daida YG, Tanaka SK, Trinacty CM, et al. Factors affecting medication adherence trajectories for patients with heart failure. Am J Manag Care. 2015;21(3):e197-e205.
41. Saccomann ICRdS, Cintra FA, Gallani MCBJ. Factors associated with beliefs about adherence to non-pharmacological treatment of patients with heart failure. Revista da Escola de Enfermagem da USP. 2014;48(1):18-24.
42. Mase R, Halasyamani L, Choi H, Heisler M. Who signs up for and engages in a peer support heart failure self-management intervention. The Journal of cardiovascular nursing. 2015;30(4 0 1):S35.
43. Siabani S, Driscoll T, Davidson PM, Najafi F, Jenkins MC, Leeder SR. Self-care and its predictors in patients with chronic heart failure in western Iran. Journal of Cardiovascular Nursing. 2016;31(1):22-30.
44. Chen AM, Yehle KS, Albert NM, Ferraro KF, Mason HL, Murawski MM, et al. Relationships between health literacy and heart failure knowledge, self-efficacy, and self-care adherence. Research in Social and Administrative Pharmacy. 2014;10(2):378-86.
45. Brown MT, Bussell J, Dutta S, Davis K, Strong S, Mathew S. Medication adherence: truth and consequences. The American journal of the medical sciences. 2016;351(4):387-99.
46. Oosterom-Calo R, Van Ballegooijen A, Terwee C, Te Velde S, Brouwer I, Jaarsma T, et al. Determinants of adherence to heart failure medication: a systematic literature review. Heart failure reviews. 2013;18(4):409-27.
47. Calvin JE, Shanbhag S, Avery E, Kane J, Richardson D, Powell L. Adherence to evidence-based guidelines for heart failure in physicians and their patients: lessons from the Heart Failure Adherence Retention Trial (HART). Congestive heart failure. 2012;18(2):73-8.
48. Davis KK, Himmelfarb CRD, Szanton SL, Hayat MJ, Allen JK. Predictors of heart failure self-care in patients who screened positive for mild cognitive impairment. Journal of Cardiovascular Nursing. 2015;30(2):152-60.
49. Wu J-R, Lennie TA, De Jong MJ, Frazier SK, Heo S, Chung ML, et al. Medication adherence is a mediator of the relationship between ethnicity and event-free survival in patients with heart failure. Journal of cardiac failure. 2010;16(2):142-9.
50. Zhang Y, Baik SH. Race/ethnicity, disability, and medication adherence among Medicare beneficiaries with heart failure. Journal of general internal medicine. 2014;29(4):602-7.
51. Buja A, Solinas G, Visca M, Federico B, Gini R, Baldo V, et al. Prevalence of heart failure and adherence to process indicators: Which socio-demographic determinants are involved? International journal of environmental research and public health. 2016;13(2):238.
52. Wu J-R, Lennie TA, Dekker RL, Biddle MJ, Moser DK. Medication adherence, depressive symptoms, and cardiac event-free survival in patients with heart failure. Journal of cardiac failure. 2013;19(5):317-24.
53. Despaigne RM, Sánchez CMD, Sarfraz A. Therapeutic adherence in outpatients with heart failure registered with a community pharmacy. Brazilian Journal of Pharmaceutical Sciences. 2012;48(1):61-8.
54. Nieuwenhuis MM, van der Wal MH, Jaarsma T. The body of knowledge on compliance in heart failure patients: we are not there yet. Journal of Cardiovascular Nursing. 2011;26(1):21-8.
55. Knafelz GJ, Riegel B. What puts heart failure patients at risk for poor medication adherence? Patient preference and adherence. 2014;8:1007.
56. Basuray A, Dolansky M, Josephson R, Sattar A, Grady EM, Vehovec A, et al. Dietary sodium adherence is poor in chronic heart failure patients. Journal of cardiac failure. 2015;21(4):323-9.
57. Aggarwal B, Liao M, Allegrante JP, Mosca L. Low social support level is associated with non-adherence to diet at 1 year in the Family Intervention Trial for Heart Health (FIT Heart). Journal of nutrition education and behavior. 2010;42(6):380-8.
58. Gravely S, Tamim H, Smith J, Daly T, Grace SL. Non-Symptom-Related Factors Contributing to Delay in Seeking Medical Care by Patients With Heart Failure: A Narrative Review. Journal of cardiac failure. 2011;17(9):779-87.
59. Thomas JR, Clark AM. Women with heart failure are at high psychosocial risk: a systematic review of how sex and gender influence heart failure self-care. Cardiology research and practice. 2011;2011.
60. Trivedi RB, Piette J, Fihn SD, Edelman D. Examining the interrelatedness of patient and spousal stress in heart failure: conceptual model and pilot data. Journal of Cardiovascular Nursing. 2012;27(1):24-32.
61. Moser DK, Watkins JF. Conceptualizing self-care in heart failure: a life course model of patient characteristics. Journal of cardiovascular nursing. 2008;23(3):205-18.
62. Son Y-J, Kim H-G, Kim E-H, Choi S, Lee S-K. Application of support vector machine for prediction of medication adherence in heart failure patients. Healthcare informatics research. 2010;16(4):253-9.
63. Heo S, Moser DK, Lennie TA, Payne-Emerson H, Welch JL, Weaver M. Development and testing of the feasibility and acceptability of a tailored dietary intervention in patients with heart failure. Journal of Cardiovascular Nursing. 2015;30(3):213-21.
64. Conraads VM, Deaton C, Piotrowicz E, Santaularia N, Tierney S, Piepoli MF, et al. Adherence of heart failure patients to exercise: barriers and possible solutions: a position statement of the Study Group on Exercise Training in Heart Failure of the Heart Failure Association of the European Society of Cardiology. European journal of heart failure. 2012;14(5):451-8.
65. Cajita MI, Cajita TR, Han H-R. Health literacy and heart failure: a systematic review. The Journal of cardiovascular nursing. 2016;31(2):121.
66. Noureldin M, Plake KS, Morrow DG, Tu W, Wu J, Murray MD. Effect of health literacy on drug adherence in patients with heart failure. Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy. 2012;32(9):819-26.
67. Chen AM, Yehle KS, Plake KS, Murawski MM, Mason HL. Health literacy and self-care of patients with heart failure. The Journal of cardiovascular nursing. 2011;26(6):446.
68. Paradis V, Cossette S, Frasure-Smith N, Heppell S, Guertin M-C. The efficacy of a motivational nursing intervention based on the stages of change on self-care in heart failure patients. Journal of Cardiovascular Nursing. 2010;25(2):130-41.
69. Maeda U, Shen B-J, Schwarz ER, Farrell KA, Mallon S. Self-efficacy mediates the associations of social support and depression with treatment adherence in heart failure patients. International journal of behavioral

- medicine. 2013;20(1):88-96.
70. Zhang KM, Dindoff K, Arnold JMO, Lane J, Swartzman LC. What matters to patients with heart failure? The influence of non-health-related goals on patient adherence to self-care management. *Patient education and counseling*. 2015;98(8):927-34.
 71. Kessing D, Denollet J, Widdershoven J, Kupper N. Psychological determinants of heart failure self-care: systematic review and meta-analysis. *Psychosomatic medicine*. 2016;78(4):412-31.
 72. Harkness K, Spaling MA, Currie K, Strachan PH, Clark AM. A systematic review of patient heart failure self-care strategies. *Journal of Cardiovascular Nursing*. 2015;30(2):121-35.
 73. van der Wal MH, Jaarsma T, Moser DK, van Gilst WH, van Veldhuisen DJ. Qualitative examination of compliance in heart failure patients in The Netherlands. *Heart & Lung: The Journal of Acute and Critical Care*. 2010;39(2):121-30.
 74. Aggarwal B, Pender A, Mosca L, Mochari-Greenberger H. Factors associated with medication adherence among heart failure patients and their caregivers. *Journal of nursing education and practice*. 2015;5(3):22.
 75. Wu J-R, Moser DK. Type D personality predicts poor medication adherence in patients with heart failure in the USA. *International journal of behavioral medicine*. 2014;21(5):833-42.
 76. Wu J-R, Song EK, Moser DK. Type D personality, self-efficacy, and medication adherence in patients with heart failure—A mediation analysis. *Heart & Lung: The Journal of Acute and Critical Care*. 2015;44(4):276-81.
 77. Farrell K, Shen B-J, Mallon S, Penedo FJ, Antoni MH. Utility of the Millon Behavioral Medicine Diagnostic to predict medication adherence in patients diagnosed with heart failure. *Journal of clinical psychology in medical settings*. 2011;18(1):1-12.
 78. Chen AM, Yehle KS, Albert NM, Ferraro KF, Mason HL, Murawski MM, et al. Health literacy influences heart failure knowledge attainment but not self-efficacy for self-care or adherence to self-care over time. *Nursing research and practice*. 2013;2013.
 79. Cameron J, Worrall Carter L, Page K, Riegel B, Lo SK, Stewart S. Does cognitive impairment predict poor self care in patients with heart failure? *European Journal of Heart Failure*. 2010;12(5):508-15.
 80. Uijen AA, Bosch M, van den Bosch WJ, Bor H, Wensing M, Schers HJ. Heart failure patients' experiences with continuity of care and its relation to medication adherence: a cross-sectional study. *BMC family practice*. 2012;13(1):86.
 81. Viana M, Laszczynska O, Mendes S, Friões F, Lourenço P, Bettencourt P, et al. Medication adherence to specific drug classes in chronic heart failure. *Journal of Managed Care Pharmacy*. 2014;20(10):1018-26.
 82. Sueta CA, Rodgers JE, Chang PP, Zhou L, Thudium EM, Kucharska-Newton AM, et al. Medication adherence based on part D claims for patients with heart failure after hospitalization (from the Atherosclerosis Risk in Communities Study). *The American journal of cardiology*. 2015;116(3):413-9.
 83. Stamp KD. Women with heart failure: do they require a special approach for improving adherence to self-care? *Current heart failure reports*. 2014;11(3):307-13.
 84. Byszewski A, Azad N, Molnar FJ, Amos S. Clinical pathways: adherence issues in complex older female patients with heart failure (HF). *Archives of gerontology and geriatrics*. 2010;50(2):165-70.
 85. Kessing D, Pelle AJ, Kupper N, Szabó BM, Denollet J. Positive affect, anhedonia, and compliance with self-care in patients with chronic heart failure. *Journal of psychosomatic research*. 2014;77(4):296-301.
 86. Alosco ML, Spitznagel MB, Van Dulmen M, Raz N, Cohen R, Sweet LH, et al. Cognitive function and treatment adherence in older adults with heart failure. *Psychosomatic Medicine*. 2012;74(9):965.
 87. Silva AFd, Cavalcanti ACD, Malta M, Arruda CS, Gandin T, Fé Ad, et al. Treatment adherence in heart failure patients followed up by nurses in two specialized clinics. *Revista latino-americana de enfermagem*. 2015;23(5):888-94.
 88. Navidian A, Yaghoubinia F, Ganjali A, Khoshsimae S. The effect of self-care education on the awareness, attitude, and adherence to self-care behaviors in hospitalized patients due to heart failure with and without depression. *PloS one*. 2015;10(6):e0130973.
 89. Riegel B, Moelter ST, Ratcliffe SJ, Pressler SJ, De Geest S, Potashnik S, et al. Excessive daytime sleepiness is associated with poor medication adherence in adults with heart failure. *Journal of cardiac failure*. 2011;17(4):340-8.
 90. Hawkins LA, Kilian S, Firek A, Kashner TM, Firek CJ, Silvet H. Cognitive impairment and medication adherence in outpatients with heart failure. *Heart & Lung: The Journal of Acute and Critical Care*. 2012;41(6):572-82.
 91. Currie K, Rideout A, Lindsay G, Harkness K. The association between mild cognitive impairment and self-care in adults with chronic heart failure: a systematic review and narrative synthesis. *Journal of Cardiovascular Nursing*. 2015;30(5):382-93.
 92. Tang H-Y, Sayers SL, Weissinger G, Riegel B. The role of depression in medication adherence among heart failure patients. *Clinical nursing research*. 2014;23(3):231-44.
 93. Riegel B, Dickson VV, Goldberg LR, Deatrick JA. Factors associated with the development of expertise in heart failure self-care. *Nursing research*. 2007;56(4):235-43.
 94. Cholowski K, Cantwell R. Predictors of medication compliance among older heart failure patients. *International journal of older people nursing*. 2007;2(4):250-62.
 95. Wu J-R, Moser DK, Chung ML, Lennie TA. Predictors of medication adherence using a multidimensional adherence model in patients with heart failure. *Journal of cardiac failure*. 2008;14(7):603-14.
 96. Meleis AI. *Theoretical nursing: Development and progress*: Lippincott Williams & Wilkins; 2012.
 97. Lazarus RS, Folkman S. *Stress, appraisal, and coping*. New York: Springer; 1984.
 98. Thomas CM. The influence of self-concept on adherence to recommended health regimens in adults with heart failure. *Journal of Cardiovascular Nursing*. 2007;22(5):405-16.
 99. Zhan L. Cognitive adaptation and self-consistency in hearing-impaired older persons: Testing Roy's adaptation model. *Nursing Science Quarterly*. 2000;13(2):158-65.
 100. Wang S-P, Lin L-C, Lee C-M, Wu S-C. Effectiveness of a self-care program in improving symptom distress and quality of life in congestive heart failure patients: a preliminary study. *Journal of Nursing Research*. 2011;19(4):257-66.
 101. Dickson VV, McCarthy MM, Howe A, Schipper J, Katz SM. Sociocultural influences on heart failure self-care among an ethnic minority black population. *Journal of Cardiovascular Nursing*. 2013;28(2):111-8.
 102. Alvarez JS, Goldraich LA, Nunes AH, Zandavalli MCB, Zandavalli RB, Belli KC, et al. Association between spirituality and adherence to management in outpatients with heart failure. *Arquivos brasileiros de car-*

- diologia. 2016;106(6):491-501.
103. Naghi JJ, Philip KJ, Phan A, Cleenewerck L, Schwarz ER. The effects of spirituality and religion on outcomes in patients with chronic heart failure. *Journal of religion and health*. 2012;51(4):1124-36.
 104. Turrise S. Illness representations, treatment beliefs, medication adherence, and 30-day hospital readmission in adults with chronic heart failure: a prospective correlational study. *Journal of Cardiovascular Nursing*. 2016;31(3):245-54.
 105. Albert NM, Levy P, Langlois E, Nutter B, Yang D, Kumar VA, et al. Heart failure beliefs and self-care adherence while being treated in an emergency department. *The Journal of emergency medicine*. 2014;46(1):122-9.
 106. Cooper LB, Mentz RJ, Sun J-L, Schulte PJ, Fleg JL, Cooper LS, et al. Psychosocial Factors, Exercise Adherence, and Outcomes in Heart Failure Patients: Insights from HF-ACTION. *Circulation Heart failure*. 2015;8(6):1044.
 107. Molloy GJ, Gao C, Johnston DW, Johnston M, Witham MD, Struthers AD, et al. Adherence to angiotensin-converting-enzyme inhibitors and illness beliefs in older heart failure patients. *European journal of heart failure*. 2009;11(7):715-20.
 108. Jeon Y-H, Kraus SG, Jowsey T, Glasgow NJ. The experience of living with chronic heart failure: a narrative review of qualitative studies. *BMC health services research*. 2010;10(1):77.
 109. Dunlay SM, Manemann SM, Chamberlain AM, Cheville AL, Jiang R, Weston SA, et al. Activities of daily living and outcomes in heart failure. *Circulation: Heart Failure*. 2015: CIRCHEARTFAILURE. 114.001542.
 110. Mohammadi E, Khoshab H, Kazemnejad A. Activities of daily living for patients with chronic heart failure: a partnership care model evaluation. *Applied Nursing Research*. 2016;30:261-7.
 111. Riegel B, Dickson VV. A situation-specific theory of heart failure self-care. *Journal of cardiovascular Nursing*. 2008;23(3):190-6.
 112. Chung ML, Lennie TA, Mudd-Martin G, Moser DK. Adherence to the low sodium diet in patients with heart failure is best when family members also follow the diet: A Multicenter Observational Study. *The Journal of cardiovascular nursing*. 2015;30(1):44.



www.revhipertension.com
 www.revdiabetes.com
 www.revsindrome.com
 www.revistaavft.com

Indices y Bases de Datos:

AVFT está incluida en las bases de datos de publicaciones científicas en salud:

OPEN JOURNAL SYSTEMS

REDALYC (Red de Revistas Científicas de América Latina, el Caribe, España y Portugal)

SCOPUS de Excerpta Medica

GOOGLE SCHOLAR

Scielo

BIREME (Centro Latinoamericano y del Caribe de Información en Ciencias de la Salud)

LATINDEX (Sistema Regional de Información en Línea para Revistas Científicas de América Latina, el Caribe, España y Portugal)

Índice de Revistas Latinoamericanas en Ciencias (Universidad Nacional Autónoma de México)

LIVECS (Literatura Venezolana de Ciencias de la Salud)

LILACS (Literatura Latinoamericana y del Caribe en Ciencias de la Salud)

PERIÓDICA (Índices de Revistas Latinoamericanas en Ciencias)

REVENCYT (Índice y Biblioteca Electrónica de Revistas Venezolanas de Ciencias y Tecnología)

SABER - UCV

EBSCO Publishing

PROQUEST