



## Artículos

### ■ **Inhibición del estrés oxidativo inducido con peróxido de hidrógeno en *Saccharomyces cerevisiae* (evaluación de un extracto acuoso de *Camellia sinensis*)**

- [Introduction](#)
- [Materials and methods](#)
- [Results](#)
- [Discussion and conclusions](#)
- [References](#)

#### Franklin Pacheco-Coello

Docente-Investigador  
Universidad de Carabobo, Laboratorio de Metales Pesados y Solventes Orgánicos, Departamento de Ciencias Básicas, Laboratorio de Bioquímica Farmacológica del Instituto de Investigaciones Biomédicas (BIOMED-UC) "Dr. Francisco J. Triana Alonso"

#### Roxy Azuaje Ramírez

Universidad de Carabobo, Laboratorio de Metales Pesados y Solventes Orgánicos

#### Corymar Orosco Vargas

Universidad de Carabobo, Laboratorio de Metales Pesados y Solventes Orgánicos

#### Doralys Ramírez Azuaje

Universidad de Carabobo, Laboratorio de Metales Pesados y Solventes Orgánicos

#### **Bioanálisis**

### **Inhibición del estrés oxidativo inducido con peróxido de hidrógeno en *Saccharomyces cerevisiae* (evaluación de un extracto acuoso de *Camellia sinensis*)**

Fecha de recepción: 14/05/2021

Fecha de aceptación: 26/05/2021

La investigación tuvo como objetivo evaluar el efecto antioxidante de un extracto acuoso de té verde (*Camellia sinensis*) orgánico, al inducir un estrés oxidativo con peróxido de hidrógeno (H<sub>2</sub>O<sub>2</sub>) a *Saccharomyces cerevisiae*. Materiales y métodos: Se emplearon hojas de té verde para la obtención del extracto acuoso. Previo a la evaluación del extracto se determinó la concentración de compuestos fenólicos por el método Folin-Ciocalteu. Para la evaluación de la capacidad antioxidante, se empleó el modelo biológico de *S. cerevisiae*. Resultados: la concentración de compuestos fenólicos totales del extracto fue 216,23±2,13 mg of GAE / g PM. En la evaluación de la actividad antioxidante se observó un marcado efecto inhibitorio del estrés oxidativo en presencia de H<sub>2</sub>O<sub>2</sub>. Conclusión: el alto contenido de compuestos bioactivos presentes en *C. sinensis*, representan una alternativa en el tratamiento y prevención de enfermedades ligadas a una elevada expresión de especies reactivas de oxígeno.

**Palabras Claves:** compuestos fenólicos, antioxidante, modelo biológico, peróxido de hidrógeno.

#### **Title**

Inhibition of hydrogen peroxide induced oxidative stress in *Saccharomyces cerevisiae* (evaluation of an extract of *Camellia sinensis*)

#### **Abstract**

The research aimed to evaluate the antioxidant effect of an organic aqueous extract of green tea (*Camellia sinensis*), by inducing an oxidative stress with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) to *Saccharomyces cerevisiae*. Materials and methods: Green tea leaves were used to obtain the aqueous extract. Prior to the evaluation of the extract, the concentration of phenolic compounds was determined by the Folin-Ciocalteu method. For the evaluation of the antioxidant capacity, the biological model of *S. cerevisiae* was used. Results: The concentration of total phenolic compounds in the extract was 216.23±2.13 mg of GAE / g PM. In the evaluation of the antioxidant activity, a marked inhibitory effect of the oxidative stress in the presence of H<sub>2</sub>O<sub>2</sub> was observed. Conclusion: the high content of bioactive compounds present in *C. sinensis*, represent an alternative in the treatment and prevention of diseases linked to a high expression of reactive oxygen species.

#### **Key Word**

## Introduction

Tea (*Camellia sinensis*) is a popular, socially accepted, safe and healthy beverage that is widely consumed around the world. Tea is beneficial to prevent and cure a variety of diseases related to oxidative stress (1-3). The health benefits ascribed to the consumption of teas are thought to be associated with their high content of bioactive ingredients such as polyphenols. The latter are secondary plant metabolites and include the subclasses of flavonoids, flavones, flavonols, flavanols, isoflavones, flavanones and anthocyanidins (4). The major active flavonoids in green tea are epicatechin (EC), epicatechin gallate (ECG), epigallocatechin (EGC) and epigallocatechin gallate (EGCG) (5,6).

The structural features of green tea catechins that significantly contribute to their antioxidant action are the presence/absence of the galloyl moiety and the number and positions of the hydroxyl groups on the rings. The latter determine their ability to interact with biological matter through hydrogen bonding, or electron and hydrogen transfer processes within their antioxidant activities. In fact, the antioxidant mechanism implies hydrogen atom transfer or single electron transfer reactions, or both (7).

The objective of this research was to evaluate, the effect antioxidant in *Saccharomyces cerevisiae*, of aqueous extract organic green tea artisanal (China).

## Materials and methods

### Origin of plant material (PM)

Non-commercial organic green tea from China, stored in its package until analysis.

### Sample preparation for extraction

2 grams of plant material were weighed. This was poured into a 400 mL Beaker, to which 200 mL of distilled water previously heated to the boiling point was added. The sample was slightly stirred for 4 min and filtered using Whatman No. 4 paper (8).



**Figure 1.** Aqueous extract of *Camellia sinensis*. Agitation prior to evaluation

### Determination of total phenolics

For the determination of total phenolics, 50  $\mu\text{L}$  were mixed with 250  $\mu\text{L}$  of the Folin-Ciocalteu 1 N reagent (Analytical grade, Merck). It was left to stand for 8 minutes and then 750  $\mu\text{L}$  of 20%  $\text{Na}_2\text{CO}_3$  and 950  $\mu\text{L}$  of distilled water were added. Was incubated for 30 min at room temperature and the absorbance was read on a Genesis 20 UV/VIS spectrophotometer (Thermo Scientific, Waltham, Massachusetts, USA). A calibration curve for Gallic Acid (Sigma-Aldrich,

Germany) was prepared with concentrations of 50, 100, 200, 300, 400, 500 and 1000 ppm. The results were expressed in mg of Gallic Acid Equivalents (GAE) / g of PM (9).

### Biological effects in cell model (*S. cerevisiae*-oxidative stress with H<sub>2</sub>O<sub>2</sub>)


This essay seeks to evaluate the ability of the compounds present in the extracts to promote the growth of yeast when subjected to oxidative stress with hydrogen peroxide, for which the following experimental sequence was carried out (10): The positive control with ascorbic acid, the negative without antioxidant and a blank only with yeast and culture medium. Before starting the absorbance measurements, *S. cerevisiae* was inoculated for 24 h under agitation at 37 ° C with 8 mL of YPD culture medium (peptone 2% w / v, glucose 2% w / v), 25 µL of yeast stock (solution of *S. cerevisiae* with a cell concentration of 5.3 x 10<sup>5</sup> cells / mL), 100 µL of ascorbic acid and 80 µL of the extract of the samples. In control negative and white were inoculated only 8 mL of YPD and 25 µL of the yeast stock. After this incubation time of the yeast under the different conditions the absorbance was measured at 600 nm, this was approximately 0.240 and corresponded to the starting point of latency period of *S. cerevisiae*. Then 160 were added µL of 0.8 mM H<sub>2</sub>O<sub>2</sub> to the positive control, negative and the extract to be evaluated at different concentrations. It was incubated at 37 ° C under stirring for 8 h and readings were taken every 30 min. Before each reading they waved the glass tubes for 10 s in vortex to ensure the homogeneity of the cells in the culture medium, then 100 µL of each test was taken and then taken to plates 96 wells orbital shaking was performed before each reading 20 s and the absorbance readings were taken at 600 nm.

### Statistical analysis

Analyses were done in triplicate, and the results were expressed as means ± standard deviation (SD). Results of antioxidant activity were each subjected to analysis of variance (ANOVA).

### Results


The results obtained show that the pure (490 µg/mL, 216.23±2.13 mg of GAE / g PM) and diluted extract was able to inhibit the oxidative stress in *S. cerevisiae*, with significant difference ( $p < 0.05$ ) in relation to the positive control of Vitamin C. The latency period was similar in both the pure, diluted and positive control extracts, with an exponential growth of *S. cerevisiae* in the pure extract from 90 min (Fig 2).



**Figure 2.** Evaluation of the antioxidant capacity of the pure extract and diluted 1/2 of *C. sinensis*

### Discussion and conclusions

Experimental data indicate that tea polyphenols may offer indirect protection by activating endogenous defense systems (11). The observed effect is due to that the phenolics compounds, EC, ECG, EGC and EGCG, present in the extract organic green tea artisanal, can indirectly regulate the expression and activity of enzymes such as CAT, SOD and glutathione (12) (Fig. 2). Although it is not known with certainty what mechanisms *S. cerevisiae* carries out to promote its growth in an oxidative stress environment, it was evidenced that the compounds present in the extract are capable of stimulating the cells to follow their normal growth path, which is perhaps related to the ability of the extracts to activate or deactivate physiological mechanisms associated with the protection of the yeast or with simple reduction  $H_2O_2$  (13, 14).



**Figure 3.** Effect on the enzyme system of phenolic compounds presentes in *C. sinensis*.

Previous studies have reported that different phenolic compounds (quercetin, resveratrol, and hesperidin), are able to protect yeast cells against damage caused by different stressing agents (14,15). The antioxidant effect of tea ingestion requires more evidence to unravel the mechanism of action and the ingredients involved. Despite there being no convincing evidence from long-term intervention studies in humans, tea flavanols are still considered to be the major candidates involved in the biological activity of teas. Possible mechanisms of action, such as the induction of an endogenous redox pathway or direct effects of polyphenol metabolites, should be elucidated so that the molecules responsible for the effect can isolated and clear-cut evidence can be obtained from long-term intervention studies(17).

Natural antioxidants are an interesting alternative in view of their variety of structures and chemical interactions, as well as the numerous biological activities they can perform. Intensive research activities are currently being carried out on plant antioxidants to meet this challenge (18). The *in vitro* and *in vivo* antioxidant activities of some natural products could be consistent in certain conditions. Overall, *in vitro* antioxidant activity and phenolic profiles of tea can provide

### Conclusions

The extract of green tea showed a protective effect in the presence of an excess of  $H_2O_2$  molecules, which favours or contributes to the metabolism of *S. cerevisiae*. The results in this research evidence that the consumption of green tea has beneficial effects on health and prevention of chronic pathologies, such as cardiovascular disease and cancer, in which oxidative stress plays a critical role.

### References

1. Tang J, Zheng JS, Fang L, Jin Y, Cai W, Li D. Tea consumption and mortality of all cancers, CVD and all causes: a meta-analysis of eighteen prospective cohort studies. Br J

- Nutr 2015; 114: 673-683.
2. Xiangbing M, Changsong G, Daiwen C, et al. Oxidative stress-induced diseases and tea polyphenols. *Oncotarget* 2017; 8, (46): 81649-81661.
  3. Peluso, I; Serafini, M. Antioxidants from black and green tea: from dietary modulation of oxidative stress to pharmacological mechanisms. *Br J Pharmacol* 2017; 174, 1195-1208
  4. Del Rio D, Rodriguez-Mateos A, Spencer JP, Tognolini M, Borges G, Crozier A. Dietary (poly) phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxid Redox Signal* 2013; 18: 1818–1892.
  5. Serafini, M; Del Rio, D; Yao, DN; Bettuzzi, S; Peluso, I. Chapter 12. Health benefits of tea. *Herbal Medicine: Biomolecular and Clinical Aspects*, 2nd ed; Boca Raton; CRC Press/Taylor & Francis, 2011:1-12.
  6. Graham HN. Green tea composition, consumption, and polyphenol chemistry. *Prev Med* 1992; 21:334–50.
  7. Lambert JD, Elias RJ. The antioxidant and pro-oxidant activities of green tea polyphenols: a role in cancer prevention. *Arch Biochem Biophys* 2010; 501: 65-72.
  8. Horžić D, Komes D, Belščak A, et al. The composition of polyphenolics and methylxanthines in teas and herbal infusions. *Food Chem* 2019; 115, 441-448.
  9. Ayala-Zavala, JF., Silva-Espinoza, A.B., Cruz-Valenzuela, R.M., Villegas-Ochoa, M.A. Antioxidant and antifungal potential of metanol extracts of *Phenillus sp.* *Rev. iberoamer. micol* 2013; 29(3); 132-138.
  10. Stinco C, Baroni, M, Naranjo R, et al. Hydrophilic antioxidant compounds in orange juice from different fruit cultivars: Composition and antioxidant activity evaluated by chemical and cellular based (*Saccharomyces cerevisiae*) assays. *J Food Compos Anal* 2015; 37: 1-10.
  11. Rahman I, Biswas S.K, Kirkham P.A. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem Pharmacol* 2006; 72(11):1439-1452
  12. Ray P.D, Huang B.W, Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cells Signal* 2012; 24: 981-990.
  13. Valérico E, Vilares A, Campos A, Pereira P, Vasconcelos, V. Effects of microcystin-LR on *Saccharomyces cerevisiae* growth, oxidative stress and apoptosis. *Toxicol.* 2014; 90: 191
  14. Albarrán G, Mendoza E, Beltrán J. M. Influence of concentration on the radiolytic decomposition of thiamine, riboflavin, and pyridoxine in aqueous solution. *Rev Colomb Quim* 2014; 43(3): 41-48.
  15. Belinha I, Amorim M.A.L, Rodrigues P, et al. Quercetin increases oxidative stress resistance and longevity in *Saccharomyces cerevisiae*. *J Agric Food Chem* 2007; 55: 2446-2451.
  16. Dani C, Bonatto D, Salvador M, Pereira M.D, Henriques J.A.P, Eleutherio E. Antioxidant protection of resveratrol and catechin in *Saccharomyces cerevisiae*. *J Agric Food Chem* 2008; 56: 4268–4272.
  17. Zhao CN, Tang GY, Cao, S.Y, et al. Phenolic Profiles and Antioxidant Activities of 30 Tea Infusions from Green, Black, Oolong, White, Yellow and Dark Teas. *Antioxidants* 2019; 8(214): 1-14.
  18. Kasangana P, Haddad P, Stevanovic T. Study of Polyphenol Content and Antioxidant Capacity of *Myrianthus Arboreus* (Cecropiaceae) Root Bark Extracts, *Antioxidants* 2015 4(2); 410-426.
  19. Tang G.Y, Zhao C.N, Xu, X.Y, et al. Phytochemical composition and antioxidant capacity of 30 Chinese teas. *Antioxidants* 2019; 8(6): 180.

**NOTA:** Toda la información que se brinda en este artículo es de carácter investigativo y con fines académicos y de actualización para estudiantes y profesionales de la salud. En ningún caso es de carácter general ni sustituye el asesoramiento de un médico. Ante cualquier duda que pueda tener sobre su estado de salud, consulte con su médico o especialista.